

# Cross-shore transport and eddies promote large scale response to urban eutrophication

Fayçal Kessouri<sup>1,2,\*</sup>, Martha Sutula<sup>1</sup>, Daniele Bianchi<sup>2</sup>, Minna Ho<sup>1,2</sup>, Pierre Damien<sup>2</sup>, James C McWilliams<sup>2</sup>, Christina Frieder<sup>1</sup>, Lionel Renault<sup>3</sup>, Hartmut Frenzel<sup>4,6</sup>, Karen McLaughlin<sup>1</sup>, and Curtis Deutsch<sup>5</sup>

<sup>1</sup>Southern California Coastal Water Research Project, Costa Mesa, CA 92626, USA

<sup>2</sup>University of California Los Angeles, Los Angeles, CA 90095, USA

<sup>3</sup>Laboratoire d'Études en Géophysique et Océanographie Spatiale, Toulouse, France

<sup>4</sup>School of Oceanography, Washington, Seattle, USA

<sup>5</sup>Princeton University, Princeton, NH 08544, USA

<sup>6</sup>Now at CICOES, University of Washington and NOAA PMEL, Seattle, USA

\*faycalk@sccwrp.org

## ABSTRACT

A key control on the magnitude of coastal eutrophication is the degree to which currents quickly transport nitrogen derived from human sources away from the coast to the open ocean before eutrophication develops. In the Southern California Bight (SCB), an upwelling-dominated eastern boundary current ecosystem, anthropogenic nitrogen inputs increase algal productivity and cause subsurface acidification and oxygen ( $O_2$ ) loss along the coast. However, the extent of anthropogenic influence on eutrophication beyond the coastal band, and the physical transport mechanisms and biogeochemical processes responsible for these effects, remain poorly understood. Here, we use a submesoscale-resolving numerical model to document the detailed biogeochemical mass balance of nitrogen, carbon and oxygen, their physical transport, and effects on offshore habitats. Despite 20 years since the last assessment and the management of terrestrial nutrients that has occurred since that time, the coastal eutrophication is still persisting. Anthropogenic nutrient inputs promote an increase in productivity, remineralization and respiration offshore, with recurrent  $O_2$  loss and pH decline in a region located 30 – 90 km from the mainland. Over 2013 to 2017, the 5-year Bight-wide spatially averaged rate of loss was  $1.3 \text{ mmol m}^{-3} O_2$ , with some locations losing on average up to  $14.2 \text{ mmol m}^{-3} O_2$ . This phenomenon persists 4 to 6 months of the year over an area of  $278,400 \text{ km}^2$  (~30% of SCB area). These recurrent features of acidification and oxygen loss are associated with eddy cross-shore transport of nutrients and plankton biomass, and their accumulation and retention within persistent eddies offshore within the SCB.

## Introduction

Rising atmospheric carbon dioxide ( $CO_2$ ) concentrations are driving global trends of warming, acidification, and oxygen ( $O_2$ ) loss in the ocean, with severe consequences for marine ecosystems. In coastal regions, these global drivers can combine with the effects of eutrophication to exacerbate declining pH and  $O_2$ , particularly in shallow and enclosed bodies of water (e.g.,<sup>1,2,3</sup>). In eastern boundary current ecosystems, wind-driven upwelling drives high biological productivity, routinely exposing shelf waters to low-pH and low- $O_2$  conditions<sup>4,5</sup>, which would suggest a minor role for coastal eutrophication in exacerbating these stressors<sup>6</sup>. However, in the Southern California Bight (SCB), an open embayment in the upwelling-dominated southern California Current System,<sup>7,8</sup> found that the effect of anthropogenically-enhanced nutrient loads from a coastal population of 23 million people is amplifying primary productivity and subsurface respiration within a 15 km band along the coast, exacerbating acidification and oxygen loss on a local scale at a rate approaching global rate that have occurred due to climate change since the pre-industrial period. However, much work remains to better understand the full spatial footprint of the influence of anthropogenic inputs on primary productivity, pH and  $O_2$  changes in the SCB, and the physical transport mechanisms responsible for spatial and temporal variability in these effects.

In open embayments such as the SCB, a key control on the magnitude of eutrophication is the degree to which the vigorous currents quickly transport nitrogen away from the coast to the open ocean before eutrophication conditions manifest<sup>6</sup>. Cross-shelf transport and entrapment of nitrogen and newly produced organic matter in persistent eddies represent a physical mechanism, antagonistic to mean currents, that can accumulate organic matter, exacerbating  $O_2$  and pH declines. Previous work has shown that mean and eddy mesoscale and submesoscale circulation features produce intense biogeochemical variability in the SCB<sup>9</sup>, made more complex by the presence of the Channel Islands<sup>10-12</sup>, abrupt topography, multiple deep basins, and a nonuniform shelf<sup>13</sup>. Offshore, the SCB circulation system is surrounded by the southward branch of the California Current and

the northward coastal current, which together generate large standing cyclonic eddies<sup>14</sup>, south and north-west of Santa Catalina Island and north of Nicolas Island. The geography of the shelf favors broad coastal-offshore exchanges, where at some places kinetic energy is more intense (e.g., Palos Verdes peninsula, and Point Dume). The island mass effect, enhanced by the wind curl, is the other feature that favors the development of intense eddy circulation in the SCB<sup>11,12</sup> and amplifies the cross-shelf exchange of energy and material. It also increases the vertical fluxes of ambient nutrients between surface and subsurface layers and favors intense seasonal phytoplankton productivity<sup>12</sup>. Nearshore, water residence time varies as a function of basin morphology and shelf and slope width, which causes spatial and temporal variability in the relative importance of mean current versus eddy cross-shelf transport.

Anthropogenic inputs of nitrogen and subsequent effects on enhanced nearshore productivity<sup>7</sup> have the potential to interact with these physical transport processes, resulting in important consequences for the rates of acidification and O<sub>2</sub> loss in a warming ocean, and the marine organisms that are exposed to them<sup>15</sup>. Point and non-point source discharges to the nearshore of the SCB are partitioned across 75 rivers and 19 ocean outfalls that discharge treated municipal wastewater effluent totaling, on average, 8 million m<sup>3</sup> d<sup>-1</sup> of nitrogen-enriched water to the SCB<sup>16</sup>. Nitrogen in river plumes discharges directly to the euphotic zone, while deep ocean outfalls are designed to achieve greater dispersion of contaminants below the euphotic zone<sup>8</sup>. Therefore, physical processes that control water column mixing, and the interaction of these riverine and ocean outfall plumes with mean currents and the eddies and jets that enhance cross-shelf transport, are key to the environmental fate and effects of the anthropogenic nutrients being discharged.

Biogeochemical mass balances analyses are routinely used in coastal numerical modeling to investigate the mechanisms responsible for the spatial and temporal changes in the sources, sinks, and transformations of nutrients, carbon, and O<sub>2</sub>, using principles of mass conservation (e.g.,<sup>17</sup>). Movement between the control volumes used for mass balance reflects physical transport (advection or vertical diffusion), which in essence link the environmental drivers with eutrophication outcomes and its biological consequences. Patterns of change in the mass balance have significant practical importance as they provide coastal managers direct evidence that link the environmental drivers to their effects, especially when those drivers result in effects that occur far away from their point of origin, as is typically the case with eutrophication<sup>18</sup>.

In this study, we assess the fate and consequences of anthropogenic nitrogen loading to the SCB on coastal nitrogen, O<sub>2</sub>, and carbon budgets, utilizing a series of simulations with a regional submesoscale-resolving physical-biogeochemical model (Materials and Methods), updated from previous work<sup>7</sup> to represent recent conditions (August 2012 to November 2017). Our analysis assesses the change between two simulations. A control simulation (CTRL) does not include any local terrestrial inputs but represents the large-scale effects of the global atmospheric CO<sub>2</sub> increase (warming and acidification). The anthropogenic simulation (ANTH) includes terrestrial inputs of nutrients (nitrogen, phosphorus, silicon, and iron), organic and inorganic carbon (C), and alkalinity from rivers, wastewater outfalls, and atmospheric deposition (Materials and Methods). When the CTRL simulation is subtracted from the ANTH simulation, we attribute the net change to these terrestrial inputs, which represent both natural and anthropogenic sources, of which >95% total N exports originate from point sources of treated municipal wastewater effluent, with a minor fraction from non-point sources<sup>8,19</sup>. We employ biogeochemical mass balance analyses to investigate the magnitude, spatial extent, and physical processes responsible for changes in patterns of net primary production (NPP), respiration, and remineralization, both inshore and offshore. We then quantify the contribution of these changes to subsurface acidification and O<sub>2</sub> loss, the basis for subsequent investigations of the effects of eutrophication on loss of suitable habitat for aerobic and calcifying taxa (Frieder et al., this issue).

The model is run at a horizontal resolution of 0.3 km over the SCB, and is forced by realistic atmospheric fields and oceanic boundary conditions, and a synoptic reconstruction of natural, non-point-source, and point-source river and ocean outfall discharges<sup>19</sup>. This configuration is part of a suite of “nested” simulations used to propagate basin-scale circulation patterns to the coastal scales where the dispersal and biological utilization of anthropogenic nutrients takes place<sup>8,20</sup>. A model skill assessment in which predictions were compared with observations demonstrated that the model captures the main distribution and variability of physical and biogeochemical variables in the region with a high degree of fidelity, including nitrogen, O<sub>2</sub>, and carbon system parameters<sup>8</sup>.

## Results

### Effects of Land-Based Nutrient Inputs on Phytoplankton Biomass, Oxygen, and pH

Bight-wide, the 5-year annually averaged (2013-2017) phytoplankton biomass shows a mean increase of 7 mmol C m<sup>-2</sup>, with values up to 154.4 mmol C m<sup>-2</sup>, relative to the CTRL scenario. Spatially, prominent increases in phytoplankton biomass occur in the coastal band, with more diffuse but ubiquitous effects in the offshore domain, where the increase reaches an average 5 mmol C m<sup>-2</sup> and up to on average 32.8 mmol C m<sup>-2</sup> over the 5 years (Fig 1A). This increased biomass occurs throughout the year, with no apparent seasonal cycle (Fig 1A bottom panel). This increase in biomass shows a strong temporal and spatial coherence with declines in subsurface oxygen and pH (Fig. 1B-C) all over the domain, with spatial averages of 1.3 mmol m<sup>-3</sup> O<sub>2</sub>, and peak losses up to 14.2 mmol m<sup>-3</sup> O<sub>2</sub>. Except within 15 km of the coast, average oxygen and pH changes show a

seasonality at the subsurface. For pH the average difference is 0.002 and a maximum difference of 0.023 occurring offshore in a 5-years average. Offshore, the subsequent pH and O<sub>2</sub> declines occur most consistently in the summer through late fall.

Nearshore, within 15 km of the coast, a previous assessment for the period 1997-2000<sup>7</sup> found that the largest anthropogenic amplifications of net primary production and biomass occurred in the San Pedro and Santa Monica Bays, where productivity increased on average by 80% and 42%, respectively. Relative to the 1997-2000 period, the more recent 2013-2017 period shows that land-based nutrients have this same persistent effect on elevating NPP and biomass along the coast (Fig. 2A), but with decreased intensity (Fig. 2B). In general, while the mean values and spatial trend have not changed much, the extremes have decreased considerably (Fig. 2C). The largest declines in productivity occurred in the San Pedro Basin, Santa Monica Bay, and nearshore of Santa Barbara. For example, the average temporal peak decreased from 700% to 167% in the San Pedro basin (Fig. 2B). Relative to the earlier simulations (1997-2000), these trends in NPP and biomass generally appear to correlate with reductions to dissolved inorganic nitrogen (DIN) export to the coast due to nutrient management (Supplemental Fig S1). Across all regions, coastal DIN export has been reduced by 13%<sup>16</sup>. Peak declines in nutrient loading have occurred in Orange County and on the San Pedro shelf, where the loads from outfalls and rivers have decreased from 83.9 to 63.4 Mg d<sup>-1</sup> (-24%<sup>19</sup>).

Offshore (bottom depth > 200m), the largest effects of land-based nutrients on biomass, oxygen, and pH during 2013-2017 occurred around south Santa Catalina Island and to a lesser extent south of the Channel Islands (Santa Rosa and Santa Cruz Islands) and northeast of San Clemente Island. The region impacted by these biogeochemical changes is approximately 278,400 km<sup>2</sup> in size. The change in the biomass is focused on the upper 50 m, corresponding to the euphotic zone, with simultaneous increases in O<sub>2</sub> and pH, signaling the effects of photosynthesis. The maximum change is generally located at the surface in the coastal band, but moves to the subsurface (~15-40 m) in the offshore domain (e.g., south of Santa Catalina Island, see the cross section in Fig 2A). Below this depth, O<sub>2</sub> and pH steeply decline, with the largest magnitude change occurring in the depth range of 40–200 m.

### Mass Balance

Eutrophication alters a series of biogeochemical processes that include photosynthesis, respiration, remineralization, and air-sea gas exchange, all of which are mediated by physical transport. These biogeochemical changes can be quantified by a mass balance analysis of DIN, organic carbon (OC), and O<sub>2</sub>. Changes in mass balance of the ANTH relative to the CTRL scenarios are calculated, focusing on the offshore domain (Fig 3).

Offshore, increased export of OC occurs to deeper waters and to the western model boundary, reflecting the amplification of both new and regenerated primary production in the offshore domain. The intensification of the vertically averaged [0-200m] photosynthesis (+3.53 mmol O<sub>2</sub> m<sup>-3</sup> d<sup>-1</sup>, +2.75 mmol C m<sup>-3</sup> d<sup>-1</sup>), relative to CTRL, is nearly balanced by increased remineralization (+2.17 mmol C m<sup>-3</sup> d<sup>-1</sup>) and respiration processes (+3.02 mmol O<sub>2</sub> m<sup>-3</sup> d<sup>-1</sup>). Air sea fluxes show an amplification of the flux of O<sub>2</sub> to the atmosphere (loss of -0.71 mmol O<sub>2</sub> m<sup>-3</sup> d<sup>-1</sup>) and the uptake of atmospheric CO<sub>2</sub> (+0.17 mmol C m<sup>-3</sup> d<sup>-1</sup>), as well as an import of O<sub>2</sub> and OC by oceanic currents (+0.2 mmol O<sub>2</sub> m<sup>-3</sup> d<sup>-1</sup>, +0.41 mmol C m<sup>-3</sup> d<sup>-1</sup>).

Changes in five-year averaged phytoplankton DIN uptake, respiration, and remineralization, integrated over 0-200 m, illustrate the spatial variability in these rates (Fig. 4A-C), which directly correspond to changes in biomass, O<sub>2</sub>, and pH (Fig. 1A-C). All along the coast on the shelf (depth < 0-200 m), DIN uptake increases by 95%, and respiration and remineralization by 50%.

In the offshore domain, the largest amplification of phytoplankton DIN uptake, respiration, and remineralization occurs southwest and southeast of Santa Catalina Island, parallel to the coast up to the southern coast of the Channel Islands (Santa Rosa and Santa Cruz) and northeast San Clemente Island, as well as northeast San Nicolas Island (Fig. 4A-C). A strong spatial coherence can be seen in the three rates with changes in biomass, O<sub>2</sub>, and pH (Fig. 1A-C). These changes are not necessarily only caused by physical transport of coastal water masses only — local biogeochemical processes can also be a factor. Seasonal climatologies of DIN uptake and respiration on the shelf and offshore reproduce this synoptic variability, but coastal remineralization shows a seasonal variability, which signals a decoupling of coastal production from the locale of its remineralization.

This decoupling is also evident in the spatial patterns of NH<sub>4</sub><sup>+</sup> uptake (Fig. 5A), which functions as a tracer for anthropogenic nitrogen inputs, versus recycled NH<sub>4</sub><sup>+</sup> uptake, which is the by-product of remineralization. NH<sub>4</sub><sup>+</sup> uptake shows a broader coastal intensification with increases up to 2 mmol m<sup>-3</sup> d<sup>-1</sup>, while NH<sub>4</sub><sup>+</sup> recycling shows a greater amplification offshore, with an increase by up to 0.5 mmol m<sup>-3</sup> d<sup>-1</sup> (Fig. 5C). NO<sub>3</sub><sup>-</sup> uptake shows variable changes along the coast (Fig. 5B), with reduction in selected nearshore regions, which could reflect inhibition by greatly increased NH<sub>4</sub><sup>+</sup> uptake. Offshore, an overall increase in NO<sub>3</sub><sup>-</sup> uptake is evident. Nitrification shows an enhancement across the domain (Fig. 5D) consistent with the patterns of NH<sub>4</sub><sup>+</sup> regeneration and remineralization.

### Physical Processes Control Fate and Transport of Anthropogenic Nutrients

The relative influence of the northern currents and the persistent eddies on the spatial distribution of anthropogenically-enhanced dissolved organic nitrogen (DON) and DIN can be seen by superposing the average currents and sea surface elevation from

the ANTH simulation on the net change (ANTH-CTRL) in integrated DON and DIN (Fig. 6A-B). In the offshore domain, anthropogenically enhanced DON is widely spread over the entire SCB. The highest concentration is located inside recurrent eddies southeast of Santa Catalina Island, west of San Clemente, along the southern branch of the northward current onshore, and offshore of the Channel Islands (toward the Islands and toward the southward CCS branch). In contrast to DIN (Fig. 6B), DON (Fig. 6A) does not appear to accumulate inside the eddy of the Channel Islands.

The 5-year averaged cross-shore physical flux of nitrogen forms ( $\text{NH}_4^+$ ,  $\text{NO}_3^-$  plus  $\text{NO}_2^-$ , and organic nitrogen) from the shelf to offshore (quantified as the transport across a vertical surface that follows the 200 m isobath) reveals multiple pathways of export for each component with a distinct vertical structure (Fig. 7A-C). For instance,  $\text{NH}_4^+$  is exported mostly by transient eddies in the upper 80 m, while the cross-shore flux by mean currents is negligible (Fig. 7A).  $\text{NO}_3^-$  fluxes show a balance between export by eddies and import by the mean current (likely reflecting upwelling; Fig. 7B). DON offshore fluxes show an important influence from both the mean current and transient eddies, both of which intensify offshore at the surface (Fig. 7C).

Alongshore,  $\text{NH}_4^+$  is exported across the shelf at rates that are proportional to the magnitude of coastal export by outfalls, which are especially notable on the San Diego, Palos Verdes, and Santa Monica shelves (Fig. 8A-B). This alongshore distribution of the flux is also in agreement with an evaluation made using Lagrangian parcels (detailed analysis and results are available in SI).  $\text{NH}_4^+$  is exported mainly in deeper layers (30-80 m) during April through June when the water column is stratified, but can be exported offshore at the surface during periods of weaker stratification (July to March; Fig. 8C). The cross-shelf flux of  $\text{NH}_4^+$  by transient eddies varies regionally along the coast, and is enhanced by circulation features with intense straining associated with anticyclonic (e.g., along Santa Monica Bay) or cyclonic eddies (e.g., south San Diego coast). A detailed analysis of the transient eddies characterization is presented in SI.

## Discussion

A key control on eutrophication in coastal ecosystems is the degree to which vigorous coastal currents quickly transport nitrogen away from the coast to the open ocean before eutrophication develops. Previous work<sup>7</sup> demonstrated that, in the SCB, an upwelling-dominated eastern boundary current ecosystem, coastal anthropogenic nutrient loading results in increased algal productivity and subsurface acidification and oxygen loss within a nearshore band (< 15 km). With expanded analysis of more recent simulations, we found that subsurface acidification and  $\text{O}_2$  loss, resulting from this enhanced productivity and respiration, is also significantly affecting the offshore regions of the SCB. We document recurrent offshore “hot-spots” of subsurface acidification and  $\text{O}_2$  loss located 30 – 90 km from the mainland, which include the deep basin south and west of Santa Catalina Island, south of the Channel Islands, and north of San Nicolas Island. In total, these hot spots occupy an area of 270,800  $\text{km}^2$  on average over the five-year simulation period, with vertical pH decline and  $\text{O}_2$  losses of up to 6% persisting 4-6 months of the year. These recurrent features offshore appear to be associated with persistent cyclonic eddies that trap and retain nutrients and organic material originating from the coast, which in turn intensifies local primary production and remineralization. Transport of DIN and DON from the coastal to the offshore region occurs by a combination of mean currents and transient eddies. The existence and role of these eddies in transporting up-welled material has been documented in for the CCS<sup>21,22</sup>. The latter consist of intensely variable currents, jets, and filaments that last a few days and are generated by a combination of baroclinic and barotropic instabilities, and interaction between tides, mean currents, and topography<sup>23</sup>. These transient features detach from the main coastal northward current to export nutrient-enriched coastal waters offshore, where they can accumulate in retentive features. Mass balance and physical transport analyses demonstrate that cross-shelf transport represents an important mechanism of nutrient delivery, antagonistic to the dilutive and dispersive effect of the main currents, that can favor accumulation of organic carbon and nutrients in retentive circulation features, exacerbating acidification and  $\text{O}_2$  loss offshore. This novel finding highlights the importance of coastal circulation features in enhancing the vulnerability of eastern boundary current ecosystems to human coastal nitrogen exports<sup>16</sup>. Given that 48% of the global wastewater discharge receives no treatment<sup>24</sup>, this suggests a pressing problem that requires further investigation in global assessments of coastal acidification and  $\text{O}_2$  loss<sup>25</sup>.

Physical processes such as persistent currents and transient eddies are the principal drivers for these phenomena, acting synergistically or antagonistically to modulate the spatial and temporal patterns of anthropogenic nutrient dispersal, primary production, and respiration changes. We find that organic matter is exported cross-shelf at the surface by the main currents when the ocean is more stratified (spring and summer), but this export can occur deeper by eddies during well mixed periods, which typically occur in fall and winter. In contrast,  $\text{NH}_4^+$ , the best tracer of anthropogenic nitrogen sources in the region<sup>16</sup>, is exported mostly by transient eddies in the upper 80 m, at rates proportional to local rates of land-based nitrogen discharges, while the cross-shore flux by mean currents remains small.

The rates of decline documented in this study and that of Kessouri et al., 2021a<sup>7</sup> approach global open-ocean rates of oxygen loss and acidification since the pre-industrial period (i.e., 5  $\text{mmol O}_2 \text{ m}^{-3}$  and 0.1 pH units<sup>26,27</sup>). These local and global changes are superimposed, and both are generally enhanced in coastal waters<sup>28-31</sup>; in combination, they can lead coastal ecosystems more rapidly toward ecological tipping points. Because of temporal and vertical habitat compression in summer to

late fall, which is most severe in the 50-150 m depth range, epipelagic—dwelling life forms are the most likely to be impacted, including calcifying zooplankton (e.g., pteropods), larval life stages of echinoderms, crustaceans, and mollusks, plus a variety of fish and invertebrates that live at the limit of their aerobic habitat<sup>32-35</sup>. Even relatively small reductions in O<sub>2</sub> and pH can drive habitat compression, a topic that is further investigated in a companion study (Frieder et al. this issue).

Our analyses show a consistent, year-round amplification of NPP by approximately 200% during the 2013-2017 period, coupled with seasonal reductions of subsurface oxygen and pH up to 15 mmol m<sup>-3</sup> and 0.02, respectively. These values are lower than, but still comparable to changes in hindcasts of SCB waters for the 1997-2000 period<sup>7</sup>. Over the past two decades, total nitrogen exports have declined 8%, with subregional declines up to 83%, focused on the Los Angeles and Orange County shelf, where most of the nitrogen export to the SCB occurs. NPP shows a clear decline in this subregion, but interannual variability of ocean state<sup>36</sup> is clearly a strong modulator of the response of the system to N reductions<sup>7</sup>, which in this analysis cannot be disentangled. Idealized scenarios testing efficacy of nitrogen reduction found that during high productivity years (e.g., 1999, 2017), the efficacy of anthropogenic nitrogen reductions in changing NPP was dampened relative to low productivity years by approximately an order of magnitude (Ho et al. this issue), driven by anomalous oceanic—atmospheric interactions that lead to shallower nitracline and nutrient enrichment of the surface mixed layer<sup>37</sup>. Thus, while coastal ecosystem eutrophication responses to point sources are more efficient than to non-point sources<sup>38</sup>, considerable complexities exist in the responses of coastal ecosystems to eutrophication<sup>18,39</sup>, leading to uncertainty in the outcome of local nutrient management strategies.

Shelf (depth < 200 m) NPP increases were roughly five times those of offshore areas, but the magnitude of O<sub>2</sub> and pH loss only doubled along the shelf, leading to questions about the mechanism for this disproportionate response. Three possible processes were investigated through mass balance: 1) coastal anthropogenic nitrogen is exported cross-shelf to increase new production offshore, 2) OC (and associated nitrogen) produced inshore is exported offshore, and 3) OC accumulates and remineralizes in offshore eddies. We found evidence that all three processes are occurring and altering subsurface OC and O<sub>2</sub> cycles, but cross-shelf export of OC and entrainment by persistent eddies was largely responsible for the disproportionate response. Persistent cyclonic eddies can be disproportionately more productive than adjacent waters fueled continuously by frontal enrichment by deeper nutrients. They have previously been recognized in other regions as cores for marine production and carbon export, such as in the Sargasso Sea<sup>40</sup>, the Alboran Sea and Algerian current<sup>41,42</sup>. More importantly, the SCB persistent eddies can be isolated for a few months during which anthropogenically intensified productivity can play an important fertilizing role, amplifying carbon export and respiration in subsurface layers, and leading to cumulative O<sub>2</sub> and pH declines. While coastal waters are generally more productive, they are shallower and more ventilated. Thus, the relationship between surface productivity and subsurface dissolved inorganic carbon (DIC) increase and O<sub>2</sub> loss may be more substantially altered by transport, air sea, and sediment fluxes, and respond in a non-linear way to nutrient inputs. The Santa Barbara Channel eddy differs from this pattern: DIN appears to accumulate, but OC does not. Therefore, anthropogenic effects on subsurface pH and O<sub>2</sub> appear reduced in this region. This unique system is saturated by a continuous supply of nutrients supported by submesoscale eddies arising from a vigorous island mass effect, i.e., the interaction of winds, currents, and topography<sup>12</sup>.

Our findings augment the evidence that suggests coastal eutrophication in upwelling-dominated ecosystems can have a significant effect on acidification and oxygen loss. Furthermore, they point to two important roles of ocean circulation in driving eutrophication not only along the coastal band, but also in offshore waters. First, the coastal circulation effectively transports DIN and OC offshore along preferential pathways associated with vigorous currents and transient eddies. Second, offshore retentive features such as persistent cyclonic eddies appear to trap anthropogenic nutrients, amplifying local primary production, biomass accumulation, export, and remineralization. This amplification in turn leads to cumulative effects subsurface pH decline and O<sub>2</sub> loss localized within these features. These co-occurring stressors, in combination with marine heatwaves, are likely to result in habitat compression for a variety of marine organisms, a topic further explored in Frieder et al., (this issue).

Uncertainty exists in these predictions. Previously conducted model skill assessment for physics and biogeochemistry lends credibility to model application to these investigations<sup>8,20,43</sup>, while skill assessment for simulations of more recent periods is ongoing. Assessment of the signal-to-noise ratio of eutrophication response in seasonal and interannual variability is another measure of uncertainty. In the SCB, interannual variability of the ocean state is a strong control on the anthropogenic nitrogen loading effects on NPP. Years with naturally higher oceanic productivity with a shallow nitracline and mixed layer experience a relatively lower anthropogenic response than a non-productive year (e.g., 1998 El Nino), when the water is warmer and more stratified<sup>7</sup>. Over the period of 2013 to 2017 examined in this study, we capture years that are climatically similar, with exception of the transitional year 2013. The period is marked by a marine heatwave from 2014 to 2017, which limited mixing between deep and surface waters. Despite this interannual variability in ocean state, our analysis demonstrates consistency in the amplification of the main mass balance terms to the anthropogenic loading over multiple years. Seasonally, the anthropogenic response on NPP is stronger in summertime due to the vertical structure of the water column and to the timing of winter mixing and phytoplankton blooms. Cross-shore eddies transport of newly-formed organic matter happens after the blooms season, and subsurface anthropogenic nutrients that have been mixed into the surface in winter and spring are also transported when stratification is stronger. Additional sources of uncertainty arise from the chaotic nature of eddies. Even though the model

shows a consistent signal for the eddy transport of anthropogenic nutrients and organic matter, quantifying the uncertainty due to intrinsic variability would be useful<sup>44</sup> and is the focus of ongoing work. Model ensemble comparisons are another helpful measure of model uncertainty that are routine in climate assessment, but are not yet within our near-term capacity.

Observational programs tend to under-sample offshore regions. Thus, future work could focus on expanding observational programs beyond nearshore waters. The two scenarios upon which this work was conducted represent an all-or-none approach; however, terrestrial inputs reflect a complex mixture of point, non-point, and natural sources, responding in a non-linear fashion to variable coastal circulation. Some outfalls and rivers may have a disproportionate role in coastal eutrophication, as the location of discharge (e.g., near headlands that cause wakes<sup>10</sup>) affects mixing and dispersion by persistent currents and eddies. Further investigations are needed to understand the relative contribution of specific outfalls and rivers along the coastline, including the import of Mexican transboundary sewage flows<sup>45</sup>, the effects of which are underrepresented in this work.

Finally, work is needed to disentangle the relative contribution of natural variability and climate change, which includes downscaling and properly resolving fine-scale pH and O<sub>2</sub> variability under future climate scenarios<sup>35</sup>, ranging from high emissions to strong mitigation, to project the range of possible pH, O<sub>2</sub>, and warming stressors that coastal ecosystems will experience<sup>46</sup>. This would provide a refined understanding of the relative importance of local anthropogenic loading to acidification and O<sub>2</sub> loss, and how much time management actions could confer increased coastal resilience, given accelerating acidification and oxygen loss from climate change<sup>29,46</sup>. This type of knowledge represents a key context to further inform the role of coastal water quality management in climate change adaptation.

## Materials and Methods

We utilized the Regional Ocean Modeling System (ROMS<sup>47,48</sup>), coupled to the Biogeochemical Elemental Cycling model (BEC<sup>49</sup>), parameterized for the California Current System (CCS<sup>20,43</sup>) to simulate the effect of land-based inputs of nutrients, carbon, and iron on coastal ocean biogeochemistry and the lower trophic ecosystem over a 20-year period. We evaluated spatial and temporal changes in the mass balance of nitrogen, carbon, and oxygen and investigated the specific processes responsible for the nearshore versus offshore fate and effects.

### Oceanic Model and Configuration

A detailed description of the ROMS-BEC model setup and configuration is presented in Kessouri et al., 2021b<sup>8</sup>, but summarized here. BEC is a multi-element (C, N, P, O, iron (Fe), and silica (Si)) and multi-plankton model that includes three explicit phytoplankton functional groups (picoplankton, silicifying diatoms, and N-fixing diazotrophs), one zooplankton group, and dissolved and sinking organic detritus. Remineralization of sinking organic material follows the multi-phase mineral ballast parameterization of Armstrong et al. (2001), and sedimentary processes have also been expanded. Particulate organic matter reaching the sediment is accumulated and remineralized with a time scale of 330 days, to provide a buffer between particle deposition and nutrient release. The ecosystem is linked to a carbon system module that tracks dissolved inorganic carbon and alkalinity, and an air–sea gas exchange module based on the formulation of Wanninkhof, 1992<sup>50</sup>.

The SCB model domain, which extends from Tijuana/Mexico to Pismo Beach (U.S. Central California coast) and about 200 km offshore, is part of a one-way nested configuration. Model nests scale from a 4-km horizontal resolution configuration spanning the entire CCS<sup>20,43</sup>, to a 1 km resolution grid covering much of the California coast<sup>51</sup> (latitude < 40.25°N), to a 0.3 km grid in the SCB<sup>8</sup>, where our investigations of local anthropogenic inputs were focused<sup>7</sup>. This grid, shown in Figure 1A, is composed of 1,400 × 600 grid points, with 60  $\Sigma$ -coordinate vertical levels using the stretching function (with the following parameters:  $\theta_s = 6$ ,  $\theta_b = 3$ , and  $h_c = 250$  m) described in<sup>48</sup>. The model is run with a time step of 30 s, and outputs are saved as 1-day averages and the physical and biogeochemical diagnostics and fluxes are saved as 1-month averages.

The 1-km model is run for the period 01/1997-12/2017, after a spin-up of 4 years. Hourly surface momentum, heat, and freshwater fluxes are derived from a 6-km resolution Weather Research and Forecast (WRF,<sup>52</sup>) atmospheric simulation<sup>43</sup> using bulk formulae modified to include a wind-current coupling parameterization necessary to attain more realistic oceanic submesoscale circulation<sup>53,54</sup>. ROMS-BEC has been validated for atmospheric forcing, physics, and biogeochemistry including O<sub>2</sub>, carbonate saturation state, primary productivity, and hydrographic parameters at a West Coast-wide scale for the period of 1995-2010<sup>20</sup> and, within the SCB where anthropogenic nutrients inputs influence coastal eutrophication, at both nearshore and offshore spatial scales for the years of 1997-2001<sup>8</sup>. More information on the model setup, forcing and coast wide validation is provided in other works<sup>8,20,43</sup>.

### Model Scenarios

A set of two simulations are used to quantify the effect of land-based nutrient export on nitrogen, oxygen, and carbon cycling. A first control simulation (CTRL) does not include any terrestrial inputs and thus represents only natural ocean cycles of nutrient, carbon, and oxygen, with the effects of atmospheric deposition and global CO<sub>2</sub> superimposed. The second includes both natural oceanic cycles of nutrients that represent these same base conditions as the CTRL scenario, to which inputs from terrestrial

sources of nutrients, C and Fe are added (ANTH). These terrestrial sources, 98% of which are anthropogenic and 95% of which are point source in origin<sup>16</sup>, consist of ocean discharges of treated wastewater discharged to 19 ocean outfalls from publicly owned treatment works (POTW), and point, non-point, and natural sources in riverine surface water inputs from 75 coastal confluences.

Previous work<sup>7</sup> quantified the effects of land-based nutrient inputs during 2/1997-01/2001. In this work, we extend the model to recent periods (08/2012-11/2017). Over this period, Bight-wide wastewater volumes have declined by 33% and coastal nitrogen loads by 14%<sup>16</sup>. The coast of Los Angeles and Orange County including Santa Monica Bay is the location with the highest volume flux of nitrogen (6.78 mmol m<sup>-3</sup>d<sup>-1</sup> of total DIN, 85% of which is in the form of NH<sub>4</sub><sup>+</sup>), representing 52% of the total SCB discharges. Farther south, the San Diego coast currently receives coastal DIN exports of 4.24 mmol m<sup>-3</sup>d<sup>-1</sup>, 74% of which was NH<sub>4</sub><sup>+</sup>. Other subregions such as Ventura/Santa Barbara and south Orange County/Northern San Diego receive appreciably less DIN, with higher proportions of NO<sub>3</sub><sup>-</sup> relative to NH<sub>4</sub><sup>+</sup> (<1 mmol m<sup>-3</sup>d<sup>-1</sup>).

## Biogeochemical Mass Balance Methods

Biogeochemical mass balance analyses are routinely used to investigate the mechanisms responsible for the spatial and temporal changes in the net sources, sinks, and transformations of carbon, nitrogen, and oxygen using principles of mass conservation. Conceptually, mass balances are conducted on units of fixed volume (i.e., a “box”), from which constituents can enter (sources) or leave (losses). Because all the grid cells are adjacent in the model, sources to one box are losses from another. Movement between the boxes is driven by physical transport (advection or vertical diffusion between cells or as point sources in specific grid cells). At any given model time-point, the constituents can be stored or can be transformed through biogeochemical processes (e.g., inorganic constituents become organic via photosynthesis). “Storage” in the box, i.e., the mass of the constituent in the box, is the value reported in the model solution for that grid cell. We utilized mass balance methods to investigate the changes in eutrophication state (O<sub>2</sub>, pH) and rate (phytoplankton nitrogen uptake, NPP, respiration, and remineralization) variables resulting from terrestrial inputs and how these effects vary nearshore (depth <200 m) and offshore. These state and rate variables are also the starting point for investigations of effects of physical transport processes on coastal biogeochemistry, described in the subsequent section.

### Nitrogen Cycle

Human activities have augmented the coastal export of DIN, including NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup>, and this DIN augments primary productivity<sup>7,55</sup>. For this reason, we calculated two types of DIN mass balances, where  $S_{DIN}$  represents sources and sinks from biogeochemical processes: 1) changes in subregional DIN loading and 2) net change in the phytoplankton DIN uptake rate, which is equivalent to photosynthesis also expressed as net primary production (NPP):  $J^{DIN} = J^{NO_3^-} + J^{NH_4^+}$

$$\partial DIN / \partial t = -\nabla \cdot (u DIN) + \partial_z (k_v \partial_z DIN) + S_{DIN} \quad (1)$$

The advection term  $-\nabla \cdot (u DIN)$  is computed using the third-order upwind scheme<sup>56</sup>, and the diffusion term  $\partial_z (k_v \partial_z DIN)$  is parameterized by the KPP boundary-layer scheme<sup>57</sup>.

### Oxygen Cycle

The O<sub>2</sub> mass balance analysis includes physical transport and biogeochemical processes, where  $S_{O_2}$  represents sources and sinks from biogeochemical processes including surface air-sea flux, photosynthesis, non-grazing mortality, grazing mortality, water-column remineralization, sediment remineralization, NH<sub>4</sub><sup>+</sup> oxidation, and nitrification (Eq. A9 in Deutsch et al. 2021):

$$\partial O_2 / \partial t = -\nabla \cdot (u O_2) + \partial_z (k_v \partial_z O_2) + S_{O_2} \quad (2)$$

### DIC cycle

DIC mass balance includes physical transport and biogeochemical processes ( $S_{DIC}$ ) of air-sea flux, photosynthesis, CaCO<sub>3</sub> production, non-grazing mortality, grazing mortality, and water-column and sediment remineralization (Eq. A11 in Deutsch et al. 2021).

$$\partial DIC / \partial t = -\nabla \cdot (u DIC) + \partial_z (k_v \partial_z DIC) + S_{DIC} \quad (3)$$

We perform a change assessment from the monthly averages for the sum of the physical transport terms and for each of the biogeochemical process terms. Monthly averages of the biogeochemical rates and transport fluxes are performed between 0 and 200 m water depth of the epipelagic layer.

## Investigations of Mechanisms Controlling Fate and Transport of Anthropogenic Nitrogen

We investigated the relative importance of three possible mechanisms responsible for spatial patterns, in particular hot spots of O<sub>2</sub> and pH loss: 1) nitrogen is advected offshore, where it fuels new production, 2) nitrogen fuels new production inshore, which is then advected offshore, and 3) biomass produced or advected offshore is recycled and is regenerated to inorganic forms. An important control on these processes is the balance of DIN and organic matter following cross-shelf transport and entrainment in eddies, versus transport in the main northward current. Transport via the northward current causes export from the Bight domain, while entrainment within eddies has the potential to increase DIN residence time and accumulate organic matter within the Bight.

In order to understand the relative importance of these three processes, we conducted several analyses: We 1) quantified the mean cross-shelf fluxes of DIN and algal organic nitrogen (ON) and partitioned those fluxes into two main transport terms: a) mean currents and b) transient eddies, which are chaotic by nature<sup>58,59</sup>, but are implicated in intensifying gradients of vertical flux, productivity, respiration, and export<sup>51,60</sup>, and 2) calculated the relative importance of nitrification and NH<sub>4</sub><sup>+</sup> production from recycled organic matter. Additional methods to support the diagnostics on eddy characterization and fate and transport of coastal plumes can be found in the Supplemental material (section: Investigation on transient eddy characteristics and transport patterns).

### Mean Current Versus Transient Eddy Decomposition

We separate the total fluxes of tracers into mean and transient eddy components following a typical Reynolds decomposition (Eqn. 4), e.g., horizontal zonal advection, where  $u$  is the velocity component, and  $N$  the concentration of a biogeochemical tracer (e.g., NH<sub>4</sub><sup>+</sup>).

$$\overline{uN} = \overline{u}N + u'N' \quad (4)$$

The overbar denotes the monthly mean, and the prime the deviation from this mean, which removes long term variations. The Reynolds decomposition is applied to the physical fluxes of biogeochemical tracers along the cross-shore direction. Total fluxes,  $uN$ , are estimated online in the model and saved at daily resolution. The mean component of the fluxes,  $\overline{u}N$ , is calculated from daily mean velocities and tracer concentrations. Eddy fluxes,  $u'N'$  are calculated by difference, based on Eqn. 4. We note that "eddy" is generally a term used for any kind of mesoscale and submesoscale activity.

### Horizontal Export of DIN and ON from the shelf

We calculate the transport across a vertical section that follows the 200 m isobath, a representative of the limit of the shelf domain<sup>61</sup>, to quantify the flux from nearshore to offshore. The calculation is done at every daily averaged velocity and nutrient concentration. For visualization purposes, the alongshore profiles are smoothed using a Gaussian filter with a 20 km horizontal window. The across-shore export from the shelf to offshore is presented as positive values, and the import to the shelf is negative (e.g., upwelling). We applied this calculation to total ON, NO<sub>3</sub><sup>-</sup> plus NO<sub>2</sub><sup>-</sup>, and NH<sub>4</sub><sup>+</sup> fluxes separately. The total ON accounts for phytoplankton and zooplankton biomass plus DON.

## References

1. Cai, W.-J. *et al.* Acidification of subsurface coastal waters enhanced by eutrophication. *Nat. geoscience* **4**, 766–770 (2011).
2. Brush, M. J. *et al.* Eutrophication, harmful algae, oxygen depletion, and acidification. *Coast. Ecosyst. Transition: A Comp. Analysis North. Adriat. Chesap. Bay* 75–104 (2020).
3. Laurent, A., Fennel, K., Ko, D. S. & Lehrter, J. Climate change projected to exacerbate impacts of coastal eutrophication in the northern Gulf of Mexico. *J. Geophys. Res. Ocean.* **123**, 3408–3426 (2018).
4. Fabry, V. J., Seibel, B. A., Feely, R. A. & Orr, J. C. Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES J. Mar. Sci.* **65**, 414–432 (2008).
5. Chan, F. *et al.* Emergence of anoxia in the California current large marine ecosystem. *Science* **319**, 920–920 (2008).
6. Fennel, K. & Testa, J. M. Biogeochemical controls on coastal hypoxia. *Annu. review marine science* **11**, 105–130 (2019).
7. Kessouri, F. *et al.* Coastal eutrophication drives acidification, oxygen loss, and ecosystem change in a major oceanic upwelling system. *Proc. Natl. Acad. Sci.* **118** (2021).
8. Kessouri, F. *et al.* Configuration and validation of an oceanic physical and biogeochemical model to investigate coastal eutrophication in the Southern California Bight. *J. Adv. Model. Earth Syst.* **13**, e2020MS002296 (2021).
9. Nencioli, F., Chang, G., Twardowski, M. & Dickey, T. Optical characterization of an eddy-induced diatom bloom west of the island of hawaii. *Biogeosciences Discuss.* **6** (2009).



10. Caldeira, R. M., Marchesiello, P., Nezhin, N. P., DiGiacomo, P. M. & McWilliams, J. C. Island wakes in the Southern California Bight. *J. Geophys. Res. Ocean.* **110** (2005).
11. Dong, C. & McWilliams, J. C. A numerical study of island wakes in the Southern California Bight. *Cont. Shelf Res.* **27**, 1233–1248 (2007).
12. Kessouri, F., Renault, L., McWilliams, J. C., Damien, P. & Bianchi, D. Enhancement of oceanic eddy activity by fine-scale orographic winds drives high productivity, low oxygen, and low pH conditions in the Santa Barbara Channel. *J. Geophys. Res. Ocean.* e2022JC018947 (2022).
13. Damien, P. *et al.* Enhanced biogeochemical cycling along the US West Coast shelf. *Glob. Biogeochem. Cycles* **37**, e2022GB007572 (2023).
14. Dong, C., Idris, E. Y. & McWilliams, J. C. Circulation and multiple-scale variability in the Southern California Bight. *Prog. Oceanogr.* **82**, 168–190 (2009).
15. Gruber, N., Boyd, P. W., Frölicher, T. L. & Vogt, M. Biogeochemical extremes and compound events in the ocean. *Nature* **600**, 395–407 (2021).
16. Sutula, M. *et al.* A baseline of terrestrial freshwater and nitrogen fluxes to the Southern California Bight, USA. *Mar. Pollut. Bull.* **170**, 112669 (2021).
17. Middelburg, J. J., Soetaert, K. & Hagens, M. Ocean alkalinity, buffering and biogeochemical processes. *Rev. Geophys.* **58**, e2019RG000681 (2020).
18. Cloern, J. E. Our evolving conceptual model of the coastal eutrophication problem. *Mar. Ecology Progress Series* **210**, 223–253 (2001).
19. Sutula, M. *et al.* Dataset of terrestrial fluxes of freshwater, nutrients, carbon, and iron to the Southern California Bight, USA. *Data Brief* **35**, 106802 (2021).
20. Deutsch, C. *et al.* Biogeochemical variability in the California Current System. *Prog. Oceanogr.* 102565 (2021).
21. Nagai, T. *et al.* Dominant role of eddies and filaments in the offshore transport of carbon and nutrients in the California Current System. *J. Geophys. Res. Ocean.* **120**, 5318–5341 (2015).
22. Gruber, N. *et al.* Eddy-induced reduction of biological production in eastern boundary upwelling systems. *Nat. Geoscience* **4**, 787–792 (2011).
23. Washburn, L. & McPhee-Shaw, E. Coastal transport processes affecting inner-shelf ecosystems in the California Current System. *Oceanography* **26**, 34–43 (2013).
24. Jones, E. R., Van Vliet, M. T., Qadir, M. & Bierkens, M. F. Country-level and gridded estimates of wastewater production, collection, treatment and reuse. *Earth Syst. Sci. Data* **13**, 237–254 (2021).
25. Yamamoto, A. *et al.* Competing and accelerating effects of anthropogenic nutrient inputs on climate-driven changes in ocean carbon and oxygen cycles. *Sci. Adv.* **8**, eabl9207 (2022).
26. Orr, J. C. *et al.* Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* **437**, 681–686 (2005).
27. Schmidtko, S., Stramma, L. & Visbeck, M. Decline in global oceanic oxygen content during the past five decades. *Nature* **542**, 335–339 (2017).
28. Breitburg, D. L., Hondorp, D. W., Davias, L. A. & Diaz, R. J. Hypoxia, nitrogen, and fisheries: integrating effects across local and global landscapes. *Annu. Rev. Mar. Sci.* **1**, 329–349 (2009).
29. Doney, S. C. The growing human footprint on coastal and open-ocean biogeochemistry. *Science* **328**, 1512–1516 (2010).
30. Booth, J. *et al.* Patterns and potential drivers of declining oxygen content along the Southern California coast. *Limnol. Oceanogr.* **59**, 1127–1138 (2014).
31. Feely, R. A. *et al.* Chemical and biological impacts of ocean acidification along the west coast of North America. *Estuarine, Coast. Shelf Sci.* **183**, 260–270 (2016).
32. Bednaršek, N. *et al.* Systematic review and meta-analysis toward synthesis of thresholds of ocean acidification impacts on calcifying pteropods and interactions with warming. *Front. Mar. Sci.* **6**, 227 (2019).
33. Bednaršek, N. *et al.* Exoskeleton dissolution with mechanoreceptor damage in larval Dungeness crab related to severity of present-day ocean acidification vertical gradients. *Sci. The Total Environ.* **716**, 136610 (2020).

34. Sato, K. N. *et al.* Response of sea urchin fitness traits to environmental gradients across the Southern California oxygen minimum zone. *Front. Mar. Sci.* **5**, 258 (2018).
35. Howard, E. M. *et al.* Climate-driven aerobic habitat loss in the California Current System. *Sci. advances* **6**, eaay3188 (2020).
36. Zaba, K. D., Rudnick, D. L., Cornuelle, B. D., Gopalakrishnan, G. & Mazloff, M. R. Annual and interannual variability in the California Current System: Comparison of an ocean state estimate with a network of underwater gliders. *J. Phys. Oceanogr.* **48**, 2965–2988 (2018).
37. Jacox, M. G. *et al.* Impacts of the 2015–2016 el niño on the California Current System: Early assessment and comparison to past events. *Geophys. Res. Lett.* **43**, 7072–7080 (2016).
38. Murphy, R. R. *et al.* Nutrient improvements in Chesapeake Bay: direct effect of load reductions and implications for coastal management. *Environ. Sci. & Technol.* **56**, 260–270 (2021).
39. Krause-Jensen, D. *et al.* Imprint of climate change on pan-arctic marine vegetation. *Front. Mar. Sci.* **7**, 617324 (2020).
40. McGillicuddy Jr, D. J. *et al.* Influence of mesoscale eddies on new production in the Sargasso Sea. *Nature* **394**, 263–266 (1998).
41. Yebra, L. *et al.* Zooplankton production and carbon export flux in the western Alboran Sea gyre (SW Mediterranean). *Prog. Oceanography* **167**, 64–77 (2018).
42. Kessouri, F. *et al.* Vertical mixing effects on phytoplankton dynamics and organic carbon export in the western Mediterranean Sea. *J. Geophys. Res. Ocean.* **123**, 1647–1669 (2018).
43. Renault, L. *et al.* Evaluation of high-resolution atmospheric and oceanic simulations of the California Current System. *Prog. Oceanogr.* **195**, 102564 (2021).
44. Da, N. D. *et al.* Contributions of wind, ocean intrinsic variability, and ENSO to the interannual variability of the South Vietnam upwelling: A modeling study. *J. Geophys. Res. Ocean.* **124**, 6545–6574 (2019).
45. Gilabert-Alarcón, C., Salgado-Méndez, S. O., Daesslé, L. W., Mendoza-Espinosa, L. G. & Villada-Canela, M. Regulatory challenges for the use of reclaimed water in Mexico: a case study in Baja California. *Water* **10**, 1432 (2018).
46. Gruber, N. Warming up, turning sour, losing breath: ocean biogeochemistry under global change. *Philos. Transactions Royal Soc. A: Math. Phys. Eng. Sci.* **369**, 1980–1996 (2011).
47. Shchepetkin, A. F. & McWilliams, J. C. The regional oceanic modeling system (ROMS): a split-explicit, free-surface, topography-following-coordinate oceanic model. *Ocean. Modelling* **9**, 347–404 (2005).
48. Shchepetkin, A. F. & McWilliams, J. C. Correction and commentary for “Ocean forecasting in terrain-following coordinates: Formulation and skill assessment of the regional ocean modeling system” by Haidvogel *et al.*, *J. Comput. Phys.* **227**, pp. 3595–3624. *J. Comput. Phys.* **228**, 8985–9000 (2009).
49. Moore, J. K., Doney, S. C. & Lindsay, K. Upper ocean ecosystem dynamics and iron cycling in a global three-dimensional model. *Glob. Biogeochem. Cycles* **18** (2004).
50. Wanninkhof, R. Relationship between wind speed and gas exchange over the ocean. *J. Geophys. Res. Ocean.* **97**, 7373–7382 (1992).
51. Kessouri, F. *et al.* Submesoscale currents modulate the seasonal cycle of nutrients and productivity in the California Current System. *Glob. Biogeochem. Cycles* **34**, e2020GB006578 (2020).
52. Skamarock, W. C. & Klemp, J. B. A time-split nonhydrostatic atmospheric model for weather research and forecasting applications. *J. Comput. Phys.* **227**, 3465–3485 (2008).
53. Renault, L., Hall, A. & McWilliams, J. C. Orographic shaping of U.S. West Coast wind profiles during the upwelling season. *Clim. Dyn.* 1–17 (2016).
54. Renault, L., Masson, S., Arsouze, T., Madec, G. & McWilliams, J. C. Recipes for how to force oceanic model dynamics. *J. Adv. Model. Earth Syst.* (2020).
55. Seitzinger, S. *et al.* Global patterns of dissolved inorganic and particulate nitrogen inputs to coastal systems: Recent conditions and future projections. *Estuaries* **25**, 640–655 (2002).
56. Marchesiello, P., McWilliams, J. C. & Shchepetkin, A. Open boundary conditions for long-term integration of regional oceanic models. *Ocean. Modelling* **3**, 1–20 (2001).

57. Large, W. G., McWilliams, J. C. & Doney, S. C. Oceanic vertical mixing: A review and a model with a nonlocal boundary layer parameterization. *Rev. Geophys.* **32**, 363–403 (1994).
58. McWilliams, J. C. The nature and consequences of oceanic eddies. *Ocean. modeling an eddying regime* **177**, 5–15 (2008).
59. McWilliams, J. C. Submesoscale currents in the ocean. *Proc. Royal Soc. A: Math. Phys. Eng. Sci.* **472**, 20160117 (2016).
60. Lévy, M., Ferrari, R., Franks, P. J., Martin, A. P. & Rivière, P. Bringing physics to life at the submesoscale. *Geophys. Res. Lett.* **39** (2012).
61. Laruelle, G. G. *et al.* Global multi-scale segmentation of continental and coastal waters from the watersheds to the continental margins. *Hydrol. Earth Syst. Sci.* **17**, 2029–2051 (2013).
62. Kessouri, F. *et al.* ROMS-BEC oceanic physical and biogeochemical model code for the Southern California Current System V2020, DOI: [10.5281/zenodo.3988618](https://doi.org/10.5281/zenodo.3988618) (2020).

## Data availability

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.


## Acknowledgements

This research was supported by the California Ocean Protection Council grant C0100400NSF, C0303000, and C0831014 (administered by California SeaGrant as R/OPCOAH-1), NOAA grants NA15NOS4780186, NA18NOS4780174, and NA18NOS4780167 (Coastal Hypoxia Research Program), and the NSF grants OCE-1419323 and OCE-1419450. This work used the Expanse system at the San Diego Supercomputer Center through allocation TG-OCE170017 from the Advanced Cyber infrastructure Coordination Ecosystem: Services and Support (ACCESS) program, which is supported by National Science Foundation grants 2138259, 2138286, 2138307, 2137603, and 2138296. Additional computational resources were provided by the Hoffman2 computer cluster at the University of California Los Angeles, Institute for Digital Research and Education (IDRE). Data and code are available following the link:<sup>62</sup>.


## Author contributions statement

FK conceived and performed the simulations, designed and performed the analysis, PD contributed to the physical analysis, MS conceived and designed the analysis, DB, LR and JCW conceived and designed the analysis and developed the modeling software. CD conceived and designed the analysis and contributed modeling software, HF and MH contributed to the setup of the model. KM, CF and MH contributed to the verification of the simulation and helped performing the biogeochemical analysis. All authors contributed to the writing of the paper.


Code is available in Kessouri et al.,<sup>62</sup>. Local land-based and atmospheric data can be found in Sutula et al.,<sup>19</sup>. Due to the size of the output, model outputs are available from the authors upon request.




**Figure 1.** 5-year (2013-2017) average difference due to terrestrial inputs. In (A) vertically integrated biomass [0-100m], (B) subsurface oxygen (at 55m depth), (C) subsurface pH (at 55m depth). Upper panel shows a 2-D view of the 5-year average. Middle row shows the average change across a persistent eddy south of Santa Catalina Island (dark-yellow solid line in the upper panels; distance is measured from NE to SW). Bottom row shows climatological difference in the offshore domain (pink contour in the upper panels) for each variable, shown as a 5-year monthly average.




**Figure 2.** (A) is the ANTH-CTRL difference in NPP between the ANTH and CTRL simulations for spring and summer averaged over the 2013-2017 period. (B) Alongshore comparison of the averaged difference in NPP in the 15km coastal band for the 1997-2000 period (red line) and the 2013-2017 period (black line). For each period, the difference between the ANTH and CTRL simulations is shown. (C) Comparison of the cumulative probability distribution of the changes shown in (B).



**Figure 3.** Schematic of the anthropogenic influence (ANTH minus CTRL) of the mass balance averaged 0 to 200 m depth in the (A) shelf and offshore domain for DIN ( $\text{mmol N m}^{-3} \text{d}^{-1}$ ), (B) offshore  $\text{O}_2$ , and (C) DIC. Arrows show the physical (red) and biogeochemical (green, blue, yellow) processes. Of note, of the  $2.00 \text{ mmol N m}^{-3} \text{d}^{-1}$  added to the shelf,  $2.25 \text{ mmol N m}^{-3} \text{d}^{-1}$  is taken up by phytoplankton, which is further supported by  $1.15 \text{ mmol N m}^{-3} \text{d}^{-1}$  in remineralized N, while  $0.9 \text{ mmol N m}^{-3} \text{d}^{-1}$  is transported offshore. On average,  $2.75 \text{ mmol C m}^{-3} \text{d}^{-1}$  of plankton carbon is produced offshore, the bulk of which is remineralized/respired, yielding a net  $\text{O}_2$  and DIC loss in the upper 0-200 m.




**Figure 4.** Consequence of eutrophication on biogeochemical processes of (A) nitrogen uptake (i.e., NPP) in  $\text{mmol N m}^{-3} \text{d}^{-1}$ , (B) respiration in  $\text{mmol O}_2 \text{m}^{-3} \text{d}^{-1}$ , and (C) remineralization in  $\text{mmol C m}^{-3} \text{d}^{-1}$ . (A1, B1, C1) Upper panels are 5-year averaged maps, averaged 0-200 m. Light contours on every map highlight 25 and 75 percentiles. Lower panel represents the assessment in a climatological year with 25 and 75 percentiles for each process, (A2, B2, C2) on the shelf (depth < 200 m) and (A3, B3, C3) offshore.




**Figure 5.** 5-year mean of the effects of terrestrial inputs (ANTH minus CTRL) on 0-200 m (A) vertically averaged phytoplankton uptake of ammonium, (B) the vertically averaged phytoplankton uptake of nitrate, (C) ammonium production which includes OC remineralization, biological release, and sediment remineralization, and (D) nitrate production via nitrification process. Units are  $\text{mmol N m}^{-3} \text{d}^{-1}$ .






**Figure 6.** Map of 5-year mean surface currents superimposed on the 5-year mean of the difference between ANTH minus CTRL for (A) ON ( $\text{mmol N m}^{-2}$ ) and (B) surface elevation superposed on DIN ( $\text{mmol N m}^{-2}$ ), integrated over 0-80 m. Surface elevation and mean currents arrows highlight the position of the main currents and the recurrent eddies offshore.



**Figure 7.** A) is the change (ANTH-CTRL) of ammonium cross-shore physical flux from shelf (depth  $< 0-200$  m) to offshore (positive sign). B) is for nitrate and nitrite. C) is for ON comprising DON, phytoplankton, and zooplankton biomasses. The flux is temporally averaged over 5 years and spatially averaged along the 200 m isobath from San Diego to Santa Barbara. Units are  $\text{mmol N m}^{-2} \text{d}^{-1}$ . All fluxes are decomposed into mean and eddy components using a 1-month temporal window.



**Figure 8.** (A) Alongshore section showing the cross-shore physical total flux of ammonium at 200 m depth limit. (B) is the vertically integrated eddy flux and (C) is the seasonality of the eddy flux. Mixed season averages all the months between October and March. Stratified season averages all the months between April and September.